

TWIGAFAS XTRA

Removes bowel roundworm segments completely



For Animal Treatment Only
Keep Out of Reach of Children

Store below 30°C.
Protect from Light.
Shake well before use



Reliability & Value

TWIGAFAS-Xtra

ORAL SUSPENSION

COMPOSITION

Contains per ml:

Levamisole Hydrochloride B.P. 3.0% w/v Oxytoclozanide B.P. 6.0% w/v



Reliability & Value

TWIGA CHEMICAL INDUSTRIES LTD.

P.O. Box 30172, NAIROBI-KENYA, Tel: 020-3942000, Fax: 020-3942405
Email:- info@twiga-chem.com Website:- www.twigachemicals.com



TWIGAFAS XTRA

Fluke and Worm Drench for Cattle, Sheep and Goats. It Controls lungworms, stomach and bowel worms and adult and immature liver flukes in livestock.

COMPOSITION

Contains per ml:

Levamisole Hydrochloride B.P. 3.0% w/v

Oxyclozanide B.P. 6.0% w/v

INDICATIONS:

TWIGAFAS XTRA is a broad spectrum anthelmintic used in the treatment and control of gastrointestinal and pulmonary nematodes and adult and immature liver fluke infections in cattle, sheep and goats including: *Trichostrongylus*, *bunostomum*, *Cooperia*, *Nematodirus*, *Haemonchus*, *Ostertagia*, *Dictyocaulus* and *Fasciola* (Liverfluke) spp. The preparation clears mature flukes present in the bile ducts and tapeworm segments. It is safe in animals of all ages and can be used in pregnant and lactating animals.

TWIGAFAS XTRA should be used at all times of the year when simultaneous treatment for liver flukes, stomach and bowel roundworms, lungworms and tapeworms is required.

CONTRA-INDICATIONS:

- o **TWIGAFAS XTRA** should not be administered to animals with an impaired hepatic function.
- o Animals should not be treated simultaneously with products containing organophosphorus compounds or diethylcarbamazine citrate. Any such treatment should not take place within 14 days before or after the use of **TWIGAFAS XTRA**

DOSAGE AND ADMINISTRATION

Shake well before use. **TWIGAFAS XTRA** should be administered as an oral drench at a dose rate of 2.5ml per 10 Kg body weight in cattle and 1 ml per 4 kg body weight in sheep and goats using a dosing gun or any other appropriate measure. Careful estimation of the live weight must be made.

Dosage

DOSAGE		
	Liveweight	Dose
CATTLE	Up to 50 kg	12.5 ml
	100 kg	25.0 ml
	150 kg	37.5 ml
	200 kg	50.0 ml
	250 kg	62.5 ml
	300 kg and over	75.0 ml

(Cattle over 300kg, a further 2.5ml per additional 10kg)



DOSAGE		
	Liveweight	Dose
SHEEP AND GOATS	Up to 10 kg	2.5 ml
	20 kg	5 ml
	30 kg	7.5 ml
	40 kg	10 ml
	50 kg	12.5ml
	60kg	15 ml

**Sheep and goats over 60 kg:
A further 1 ml per every additional 4 kg



WITHDRAWAL PERIOD

- Sheep may be slaughtered for human consumption only after 15 days from the last treatment.
- Cattle may be slaughtered for human consumption only after 12 days from the last treatment.
- Milk for human consumption must not be taken during treatment
- Milk for human consumption may be taken only from cattle, from 36 hours after the last treatment
- Not suitable for use in sheep that are being milked for human consumption



CAUTION

For animal treatment only. Keep out of reach of children. Store away from direct sunlight below 30°C

AVAILABLE PACKS

125ml, 500ml and 1Litre