

# Vigo

**Plant-based Biostimulant**  
**For Vigorous Root and Shoot Growth**

*Increased Quality and  
Quantity of Yield*



**TWIGA CHEMICAL INDUSTRIES LTD.,**  
**Contact:-** Nanyuki Road - Industrial Area P.O. Box 30172, 00100 GPO Nairobi - Kenya. Tel: 3942000 /3942300, Fax: 3942405,  
E-mail: [info@twiga-chem.com](mailto:info@twiga-chem.com) Website: [www.twigachemicals.com](http://www.twigachemicals.com)

Parry patented Biostimulant, **VIGO™** is of 100% plant origin and helps the crop to achieve a vigorous root and shoot growth and enables growers to reap higher yields.

## PRODUCT FEATURES

- Get absorbed quickly by the leaves and integrates easily with plant's metabolism
- Helps in improving micro flora in the soil leading to nutrients assimilation.
- Improves absorption and transportation of nutrients inside the plant and helps in increasing nutrients use efficiency.
- Increases degree of photosynthesis in plant and make the plant lush green.
- Activates phytohormones, increases cell multiplication and thus improves crop growth and vigor.
- Facilitates Fecundation, increases fruit, pod, and seed development, increases yield and also enhances the cosmetic value of the produce.
- Strengthens plant and improves defense mechanism to fight against abiotic and biotic stresses.
- Eventually **VIGO™** usage leads to increased return on investment to the growers.

## COMPOSITION

Plant extract: 1.0% {containing vasicine 0.01%}. Other ingredient: 99.0%. Total: 100%

## RECOMMENDED CROPS

Vegetables, Fruits, Flowers, Cereals, Pulses, Oilseed crops, Spices, Cotton, Soya Beans, Sugarcane, Tea, Coffee, Cardamom etc

## DOSAGE

**2L/ha (2ml/L of water)**

### Time of application


Can be applied at all stages of crop growth- see table below for details

### For best results:

Ensure thorough coverage, repeat application on need basis at 2-3 weeks intervals.

## RECOMMEDATIONS ON USAGE:

CROPS	Dose rate	Timing of Application
Roses, carnations, other ornamentals.	2mls/litre of water 2lts/ha	Apply at any stage of crop growth, repeat 2-3 times, or on need basis.
Snow peas, Garden peas, Fine beans, runner beans	2mls/litre of water 2lts/ha	2-3 times from pre-blossoming repeat at 15 day intervals.
Broccoli, Tender stem broccoli	2mls/litre of water 2lts/ha	Apply at any stage of crop growth, repeat 2-3 times on need basis
Strawberry/raspberry	2mls/litre of water 2lts/ha	Before, during and after flowering
Chives, onions, Garlic	2mls/litre of water 2lts/ha	Every 15 days after transplanting
Tomatoes, pepper, capsicum	2mls/litre of water 2lts/ha	From transplanting till harvest starts
Cabbages, kales, spinach	2mls/litre of water 2lts/ha	Every 15days after transplanting
Potatoes	2mls/litre of water 2lts/ha	From 5 leaf stage at 2-3 week intervals
Baby corn, maize	2mls/litre of water 2lts/ha	At 3-4leaf stage, repeat 2-3 times
Grapes	2mls/litre of water 2lts/ha	Every 15 days from flower bud formation, blossoming, fruit set.
Cucurbits	2mls/litre of water 2lts/ha	2-3 times, before flowering, after flowering and at fruit setting
Coffee	2mls/litre of water 2lts/ha	Before flowering and at berry expansion.
Tea	2mls/litre of water 2lts/ha	Every 2 weeks, after pruning
Rice	2mls/litre of water 2lts/ha	At tillering stage, repeat 2-3 times.
Macadamia	2mls/litre of water 2lts/ha	2-3 applications-after harvesting and before flowering and during
Fruit trees-mangoes, citrus, avocado;	2mls/litre of water 2lts/ha	3 applications-before flowering, after flowering during and after fruit set.
Wheat and barley	2mls/litre of water 2lts/ha	At tillering stage and repeated 2-3 times.
Pineapples	2mls/litre of water 2lts/ha	2-3 times, before flowering, during fruit set and after fruit set.



## For Vigorous Root and Shoot Growth