

MAELEZO: STRETCH 20 WP ni Kiuwa kwekwe kabla hayajakua yenye kudhibiti magugu yenye majani mapana na chenye uwezo wa kuchagua na kupenya ndani ya aina nyingi za kwekwe na kuziangamiza. Huua na kuangamiza kwekwe wanao, ota haraka na changa kwenye mashamba ya ngano.

JINSI YA KUTUMIA: Isipokua kuzuia kwekwe kabla ya kupanda nyunyiza STRETCH 20 WP baada ya nafaka kuota au wakati kwekwe warehitimu centimita 5 au chini ya centimita 5 kwa urefu na upana/kipenyo.

Matokeo mazuri ni wakati wa mvua ambapo udongo una yevu kipindi cha wiki 4 baada ya kunyunyiza STRETCH 20 WP.

Usinyunyize STRETCH 20 WP kwa mimea iliyo katika hali ya ufadhako utokanao na hali ya kiangazi au kukosa maji.

Inavyofanya Kazi:

Mizizi na majani ya mimea hufyonza dawa kwa haraka huku dawa ikisonga kwenye maeneo ya mimea inayoota kwa haraka (*Tips of roots and foliage*).

Uteuzi wa dawa ni kulingana na spishi ya mimea na uwezo wa mimea huo kumetaboli metsulfuron.

Matayarisho ya dawa: Tia maji kwenye bomba la kuchanganyisha/kunyunyiza hadi kiwango cha nusu. Tia kiasi kinachohitajika cha STRETCH 20 WP kisha jaza bomba hilo kwa maji, huku ukikoroga vizuri. Nyunyiza STRETCH 20 WP wakati wa vuguvugu wa joto kwekwe/magugu yanapokuwa. Kunyunyiza wakati wa ukame hupeana matokeo duni.

VIPIMO VYA MATUMIZI

MMMEA	KWEKWE / MAGUGU	KIPIMO
Ngano	Magugu yenye matawi mapana	gramu 20/hektari (gramu 2/ Lita 20)

Tumia lita 200 - 400 za maji kwa kila hektari moja.

MADHARA KWA MIMEA: Usinyunyize dawa hii au kuiruhusu kugusa pamba, zabibu, miti ya matunda, mboga, maua na mapambo.

TAHADHARI: Vaa nguo za kujikinga, Viatu, glavu, osho (face shield) na chombo maalum cha kupumua unapatayarisha mchanganyiko wa dawa na maji na unaponyunyiza dawa. Jihadhari usipumue mvuke wa dawa, mchanganyiko wa dawa usingie kwenye macho au kumwagikia ngozi na nguo ulizovalia. Usinyunyize dawa hii ukielekea upande upepe unakotoka. Usivute sigara, usile au usinywe wakati unapotumia dawa hii.

UHIFADHI NA UTUPAJI: Hifadhi dawa hii katika chombo chake halisi ambacho kimefunikwa na kuwekwa mahali ambapo ni salama, mbali na watoto, watu wasioruhusiwa, vyakula, maji na vyakula vya mifugo, moto au joto kali. Pochi tupu la dawa yafaa kusafishwa mara tatu, na maji hayo kumwagwa kwenye bomba la mchanganyiko wa dawa. Tupa pochi tupu mahali palipo salama na paliporuhusiwa, mbali na makao ya watu na vianzo vya maji, kulingana na sheria za utupaji. Usitumie pakiti tupu kwa matumizi mengine yeyote.

MADHARA KWA MAZINGIRA: Mvuke wa STRETCH unaweza kutiririka kwenye mimea isiyotarajiwa na kuidhuru. Usichafue mazingira kwa kunyunyiza, kumwaga mabaki ya dawa, kutupa au kusunza mkoba wa dawa kwa mito, visima au vidimbwi na chemichem.

MAELEZO YA HUDUMA YA KWANZA: Ikimezwa: Mpeleke mgonjwa na kibandiko hiki kwa daktari mara moja. Usifanye mjeruhi kutapika. Usimpe mgonjwa chochote kupitia kwa mdomo ikiwa amezirai. **Ikimwagikia ngozi:** Vaa nguo zilizomwagikiwa na dawa mara moja. Oga vizuri kwa kutumia maji na sabuni kwa muda wa dakika 15-20. **Ikiingia kwenye macho:** Yasafishe kwa maji yanayotiririka kwa muda wa dakika 15. **Ikivutwa:** Mpeleke mgonjwa pahala palipo na hewa safi. Usimtapishe aliyemeza sumu. Usimtapishe mgonjwa lakini mpeleke kwa daktari mara moja.

HABARI KUHUSU SUMU: Metsulfuron-methyl ni sumu ya asili ya kemikali ya Urea.

Dalili za kusumika

- Hapana uwezekano wa kusumika kwa viungo isipokuwa wakati imemezwa kwa wingi.
- Sumu ya asili ya Urea huwasha macho, ngozi na utandu wa kamasi/pua.
- Kukohoa na ugumu wa kupumua
- Kichefuchefu, kutapika, kuhara, kuumwa na kichwa, wasiwasi
- Kuathirika kwa metaboli ya protini, mapafu na upungufu wa uzito wa mwili panapo dhara ya muda mrefu.

Tiba ya sumu: Haina tiba maalum. Tibu kulingana na dalili za sumu.

IWAPO DHARURA: Wasiliana na nambari hii isiyomalipo-0800720021/0800730030 (masaa 24)

ILANI KWA MTUMIAJI: Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyosalama.

THIBITISHO: Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambazo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima akubali dawa hii katika hali hiyo.



STRETCH[®]

HERBICIDE WETTABLE POWDER 20 WP

Post-emergence herbicide for the control of broad leaf weeds in wheat (Kiuwa kwekwe kabla ya kuchipuka, chenye kudhibiti kwekwe za majani mapana kwenye Ngano.)

Distributed by (Husambazwa na):
OSHO CHEMICAL INDUSTRIES LTD.

P.O. Box 49916, 00100-Nairobi- KENYA
PILOT Lines :- (+254) 020 3912000,
0711 045 000, 0732 167 000
E-mail: oshochem@oshochem.com
Web: www.oshochem.com

Manufacturer (Imetengenezwa na):

ATUL LIMITED
AGROCHEMICALS DIVISION
ATUL 396020, GUJARAT, INDIA



NET CONTENTS (KIASI):

1 kg



COMMERCIAL & AGRICULTURAL CLASS
SEHEMU YA BIASHARA NA KILIMO
GUARANTEE (DHAMANA): Metsulfuron Methyl 20%
REGISTRATION NO.(NAMBARI YA USAJILI): PCPB (CR) 1152
READ THE LABEL BEFORE USING (SOMA KIBANDIKO KABLA YA KUTUMIA)
KEEP LOCKED & OUT OF REACH OF CHILDREN (FUNGIA MBALI NA WATOTO)



Batch No. (Nambari ya fungu):

Mfg Date (Tarehe ya kutengenezwa):

Expiry Date (Tumia kabla ya):

Shelf Life (Maisha Rafuni): 2 years in original unopened container stored in a cool, dry & dark place (Miaka miwili toka tarehe ya kutengenezwa ikiwa kwenye chombo chake asili kichhofunguliwa kwenye sehemu tulivu na kavu.)

In case of an Emergency call Toll Free No. (24 hrs) (Wakati wa Dharura piga Simu bila malipo (masaa 24) 0800720021 / 0800730030



DESCRIPTION: STRETCH 20 WP is a systemic, selective post-emergent herbicide for the control of broad-leaved weeds in wheat when weeds are young and actively growing.

DIRECTIONS FOR USE: With the exception of pre-plant weed control, apply STRETCH 20 WP at the early post emergence stage of the crop (3 leaf stage to before booting) when weeds are not greater than 5cm in height or diameter. Best control is achieved when rainfall wets the soil to a depth of 5cm within 4 weeks of application. Do not apply to crops that are stressed by any condition.

Mode of Action: The active ingredient is absorbed by roots and foliage and rapidly moved to the growing tips of roots and foliage of plants. Selectivity is based on the plant species' capability to metabolise metsulfuron.

Preparation of Spray mixture: Fill the spray tank half full, then add the required amount of STRETCH 20 WP and continue filling the tank with the balance of water. Keep agitator running when filling the tank and during spray operations since the mixture must be kept in suspension at all times.

APPLICATION RATE

CROP	WEED	APPLICATION RATE
Wheat	Broad leaf weeds	20 gm/ha (2g/20Litres)

Spraying volume: Use 200 - 400L of water per hectare.

PHYTOTOXICITY: Do not apply this product to or otherwise permit it to come into contact with cotton, grapes, fruit trees, vegetables, flowers or ornamentals.

PRECAUTIONS: Do not smoke, eat or drink when using this product. Do not inhale the spray mist. Always wear protective clothing, boots, gloves, masks and a face shield when spraying this product.

STORAGE AND DISPOSAL: Store product in a cool, dry, locked place out of reach of children away from food or feed. Triple rinse empty container and add washings to spray mixture, then dispose empty container in a safe and authorised place away from human habitation and water sources, in accordance with national regulations and legislation. Do not reuse containers for any other purpose.

ENVIRONMENTAL HAZARD: Spray drift or runoff may adversely affect non-target plants. This product may injure cotton, beans, peas, grapes, ornamentals, etc. Do not contaminate water sources, ponds or ditches with the concentrate, its solution or empty containers.

FIRST AID MEASURES: If swallowed: Do not induce vomiting. get medical attention. Wash mouth with water and take large quantity of water. Never give anything by mouth to unconscious person. Gastric lavage is done and give the victim charcoal. Move to the hospital without delay. **Skin contact:** Remove contaminated clothing and Wash contaminated skin with plenty of clean water and soap for 15-20 minutes. **Eye contact:** Flush immediately with clean water for 15 minutes at least. **Inhalation:** Don't induce vomiting, gastric lavage is done and call a physician without delay.

TOXICOLOGICAL INFORMATION: Metsulfuron-methyl is a Urea compound.

Symptoms of Poisoning with Urea Compounds

- Systemic toxicity is unlikely unless large amounts have been ingested.
- Many substituted ureas are irritating to eyes, skin and mucous membranes.
 - Coughing and shortness of breath.
 - Nausea, vomiting, diarrhea, headache, confusion and electrolyte depletion.
 - Protein metabolism disturbances, moderate emphysema, and weight loss with chronic exposure.

Antidote: No specific antidote. Treat symptomatically.

In case of Emergency Call Toll free No. 0800720021 / 0800730030 (24 hrs)

NOTICE TO USERS: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

SELLER'S GUARANTEE (WARRANTY): Seller's guarantee is limited to the terms set out on the label and subject thereto; the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.